



ViaZen Digestion Spec Sheet

FOR HEALTH PROFESSIONALS

Presentation

90 capsules

Anticipated results

- Rapid and effective relief of dyspepsia symptoms
- Improve nutrients absorption
- Elimination of nutritional deficiencies associated with a bad digestion
- Disappearance of tiredness after the meals
- Eating become again a pleasure



Secondary anticipated results

- Gastro-intestinal mucosa integrity.
- Healthy intestinal flora.
- Immune balance.
- Better vitality.
- Eating become again a pleasure.

Who needs ViaZen Digestion?

Specifically developed for the active consumer, victim of a high-speed lifestyle and who wants to relieve its bad digestion, with a natural solution, effective, simple, user-friendly and without side effects. An optimal digestion is essential to a general good health.

Clinical symptoms of bad digestion

- Slow digestion feeling
- Heartburn
- Frequent eructation
- Stomach swelling
- Stomach heaviness


- Abdominal pain
- Hiccup
- Nausea and vomiting
- Bloating
- Gas

- Abdominal cramps
- Fatty substances in the stool
- Food fragments in the stool
- Constipation
- Diarrhea

- Headaches after eating
- Tiredness after eating
- Tiredness after eating
- Loss of appetite or early feeling of satiety

ViaZen Digestion, a powerful synergy of 4 active medicinal ingredients

The ingredients' synergy of ViaZen Digestion is the key of the effectiveness of this SUPERIOR FORMULA.



Non-medicinal ingredients

Microcrystalline cellulose, vegetable magnesium stearate, gelatine. Viazen Digestion is free of wheat, soya, corn, yeast, peanut, gluten, egg and dairy products. It does not contain preservatives, sweeteners, colouring agents, artificial flavours and is exempt of GMO.

Recommended use

Viazen Digestion is recommended in presence of discomforts associated with a bad digestion. To relieve chronic problem, Viazen Digestion will be taken on a regular basis, which means at a rate of 1 capsule with every meal. The results will appear after a few days of treatment. Viazen Digestion can also be taken in a sporadic way, to prevent bad digestion in presence of heavy meals, alcohol consumption (wine, beer, spirits and liquors) and with late meals in evening or in presence of a lot of stress. The dosage will be of 2 capsules at the time.

Be sure to ingest Viazen Digestion a few hours before or after taking other medication.

Action mechanisms

Bad digestion results from an inadequate digestive transit, which is too rapid or too slow, caused mainly by the insufficient secretion of digestive juices, including saliva, gastric juices, pancreatic enzymes, intestinal juices and the bile.

These anomalies generate intestinal fermentations and putrefactions and predispose to an intestinal dysbiosis (abnormal intestinal flora). It also reduces the absorption of nutrients and increases the risk of developing a lot of health disorders.

The **porcine pancreatine** present in Viazen Digestion contains enzymes that will optimize the digestive process.

The active ingredients of the **ginger root** will complete the action of the pancreatine by improving the mechanical portion of the digestive process. Its active ingredients including the gingerols will increase the gastro-intestinal motility and gastric emptying.

The emollient action of the **marshmallow root** will ensure a better elimination. Gastro-intestinal mucosa integrity will be ensured by different ways.

First of all, by the gastroprotective effect of the ginger; and then, by the anti-inflammatory properties of the marshmallow root on the gastro-intestinal mucosa. Lastly, by the presence of **zinc** which plays an important role in the healing process.

Marshmallow root mucilage polysaccharides will be digested by the intestinal flora bacteria and will thus act as prebiotics and will contribute to maintain a healthy intestinal flora.

Cautions

Consult a health care practitioner if symptoms persist or worsen. Consult a health care practitioner prior to use if you are pregnant or breastfeeding or if you have diabetes, pancreatitis, pancreatic exocrine insufficiency or cystic fibrosis. Do not use if you are sensitive to pork proteins or to pancreatic enzymes. Nausea, vomiting, abdominal pain/epigastric pain and/or heartburn have been known to occur, in which case discontinue use and consult a health care practitioner.

References

Gagnon R. La Nutriérapie. Collection Douce Alternative, Amyris. 2004. 287 pages. Marieb E. Anatomie et physiologie humaines. ERPI.1993. 1013 pages. Suarez F. Pancreatic Supplements Reduce Symptomatic Response of Healthy Subjects to a High Fat meal. Digestive Diseases and Sciences. 1999. Vol. 44. No. 7 1317-21. Sarnier M. Treatment of Pancreatic Exocrine Deficiency. World J. Surg. 2003. 27, 1192-5. Camilleri M. Functional dyspepsia : mechanisms of symptom generation and appropriate management of patients. Gastroenterol Clin North Am. 2007. Sept;36(3):649-64. Platel K et al. Influence of dietary spices or their active principles on digestive enzymes of small intestinal mucosa in rats. Int J Food Sci Nutr. 1996. Jan;47(1):55-9. Shukla Y et al. Cancer preventive properties of ginger: a brief review. Food Chem Toxicol. 2007. May;45(5):683-90. al-Yahya MA et al. Gastroprotective activity of ginger zingiber officinale rosc., albino rats. Am J Clin Med. 1989;17(1-2):51-6. Yamahara J et al. J Ethnopharmacol. 1988. Jul-Aug;23(2-3):299-304.

www.medicinescomplete.com / www.naturaldatabase.com / www.passeportsante.net / www.hc-sc.gc.ca/dhp-mps/prodnatur/applications/licen-prod/monograph/mono_list_f.html#m
www.mydyspepsia.com/French/InfoDyspepsia.asp / www.fr.wikipedia.org/wiki/Dyspepsie
www.naturalstandard.com/www.who.int/medicinedocs/index.fcgi?sid=zNnGhaRZ9ee80ca60000000476162bf&a=d&c=medicinedocs&d=Js4927e.3